

On the existence of smooth Cauchy steep time functions

E Minguzzi

Dipartimento di Matematica e Informatica “U. Dini”, Università degli Studi di Firenze, Via S. Marta 3, I-50139 Firenze, Italy.

E-mail: `ettore.minguzzi@unifi.it`

Abstract. A simple proof is given that every globally hyperbolic spacetime admits a smooth Cauchy steep time function. This result is useful in order to show that globally hyperbolic spacetimes can be isometrically embedded in Minkowski spacetimes and that they split as a product. The proof is based on a recent result on the differentiability of Geroch’s volume functions.

1. Introduction

Let (M, g) be a spacetime, namely a Hausdorff, paracompact, connected, time-oriented Lorentzian manifold, endowed with a (C^3) metric g of signature $(-, +, \dots, +)$. Since every C^r manifold is C^r diffeomorphic to a C^∞ manifold [6], we shall not bother with the the degree of differentiability of M , as we can suppose to have been given a smooth atlas.

On a spacetime a *time function* $\tau : M \rightarrow \mathbb{R}$ is a continuous function which increases over every causal curve, that is, $x < y \Rightarrow \tau(x) < \tau(y)$, where as usual, we write $x < y$ if there is a future-directed causal curve connecting x to y . A *Cauchy time function* is a time function for which the level sets are Cauchy hypersurfaces, namely closed acausal sets intersected exactly once by any inextendible causal curve. A time function τ is *steep* if it is continuously differentiable and $-g(\nabla\tau, \nabla\tau) \geq 1$.

The existence of smooth Cauchy time functions in globally hyperbolic spacetimes has been proved through different approaches in [1–3]. It implies that these spacetimes admit a smooth splitting as a product $\mathbb{R} \times S$, $g = -\beta^2 dt^2 \oplus h_t$, $\beta : M \rightarrow (0, +\infty)$, where t is the Cauchy time function and h_t is a time dependent Riemannian metric over the Cauchy hypersurface $\{t\} \times S$. Recently, Müller and Sánchez [7] proved that the Cauchy time function can be found steep, a fact which implies that (M, g) can be embedded in N -dimensional Minkowski spacetime for some $N \geq 2$, and that β can be chosen such that $\beta \leq 1$. This result can also be used to express a formula for the Lorentzian distance in terms of the family of time functions [8].

In this work we provide a simple proof of the existence of smooth Cauchy steep time functions in globally hyperbolic spacetimes by using some recent results on Geroch's volume functions.

Let (M, g) be a spacetime and let us consider a non-negative continuous function φ . The Geroch's volume function is

$$\tau_\varphi^\pm(p) = \int_{J^\pm(p)} \varphi d\mu_g, \quad (1)$$

where $d\mu_g$ is the volume element of g .

In a recent joint work with J. Grant and P. Chruściel [2] we proved the following lemma, which basically follows from the differentiability properties of the exponential map.

Lemma 1.1. *In globally hyperbolic spacetimes the functions $\tau_\varphi^\pm$ are continuously differentiable for all continuous compactly supported non-negative functions φ . The gradient at $p \in M$ is past-directed timelike or vanishing depending respectively on whether $E^\pm(p)$ intersects or not the set $\{q : \varphi(q) > 0\}$.*

For completeness we mention that the gradient is given by

$$\nabla_X \tau_\varphi^\pm = \int_{E^\pm(p)} \varphi J(X) d\mu_g,$$

where $J(X)$ is the Jacobi field obtained solving the Jacobi equation over each generator $\gamma(s)$ of the horismos $E^\pm(p)$, with initial condition $J(0) = X$, $(\frac{d}{ds}J)(0) = 0$.

In the proof of the main Theorem 1.3 it will be useful to keep in mind the next topological result.

Lemma 1.2. *Let K be a compact subset of a topological space and let $\{U_s\}_{s \in S}$ be a locally finite family of subsets. Then K is intersected by at most finitely many elements of the family.*

Proof. Let A_i be the open (in K) subset of K made of points which admit a neighborhood which intersects at most i elements of $\{U_s\}_{s \in S}$. Clearly, $A_i \subset A_{i+1}$, and $\bigcup_i A_i = K$. However, this open covering of K admits a finite subcovering which proves that $K = A_j$ for some j . Since every $p \in K$ admits an open neighborhood N_p which intersect at most j elements of the family $\{U_s\}_{s \in S}$, and since K admits a finite subcovering of say, r elements of the form N_{p_i} , K is intersected at most by rj elements of the family $\{U_s\}_{s \in S}$. $\square$

We are ready to prove the existence of smooth Cauchy steep time functions. The proof uses just the previous Lemmas and Geroch's topological splitting theorem.

Theorem 1.3. *Let (M, g) be a globally hyperbolic spacetime. There exists a smooth Cauchy time function $\tau : M \rightarrow \mathbb{R}$ with timelike past-directed gradient $\nabla\tau$ which is steep, namely $-g(\nabla\tau, \nabla\tau) \geq 1$.*

Proof. Just in this proof we shall say that τ is ‘steep’ if $-g(\nabla\tau, \nabla\tau) > 1$ with the strict inequality. According to Geroch’s topological splitting [4, 5] there is a continuous Cauchy time function $t: M \rightarrow \mathbb{R}$. The level sets S_t are Cauchy hypersurfaces. We are first going to construct a continuously differentiable function τ^- (resp. $-\tau^+$) over M , with past-directed timelike or vanishing gradient, which is steep over $J^+(S_0)$ (resp. $J^-(S_0)$) and such that $\tau^- - t > 0$ over $J^+(S_0)$ (resp. $-\tau^+ < t$ over $J^-(S_0)$). Then $\tau' = \tau^- - \tau^+$ will be clearly continuously differentiable, steep and Cauchy over M .

Next, by [6, Theor. 2.6] $C^\infty(M, \mathbb{R})$ is dense in $C^1(M, \mathbb{R})$ endowed with the Whitney strong topology [6, p. 35], thus we can find $\tau \in C^\infty(M, \mathbb{R})$ which approximates τ' , up to the first derivative, as accurately as we want over M . In particular, we can find τ such that $|\tau - \tau'| < 1$ and $-g(\nabla\tau, \nabla\tau) > 1$, where the former inequality implies that τ is Cauchy, and the latter inequality implies that τ is steep.

Let us construct τ^- , the plus case being analogous. For every $p \in J^+(S_0)$ there is a open neighborhood V_p with compact closure contained in $I^+(S_{-1})$. Let $V_0 = M \setminus J^+(S_0)$, by paracompactness the open covering $\{V_0\} \cup \{V_p : p \in J^+(S_0)\}$ admits a locally finite refinement (necessarily countable by Lemma 1.2 and σ -compactness of M) and a corresponding partition of unity $\{\varphi_i, i \geq 0\}$, where, defined $U_j = \{q : \varphi_j(q) > 0\}$ we have for $j \geq 1$, $\text{supp } \varphi_j = \overline{U_j} \subset I^+(S_{-1})$.

Let $K_i \subset J^+(S_0)$, be a sequence of compact sets such that $K_0 = \emptyset$, $\bigcup_i K_i = J^+(S_0)$, $K_i = J^-(K_i) \cap J^+(S_0)$ and

$$K_i \cup \bigcup_{j: U_j \cap K_i \neq \emptyset} \overline{U_j} \subset K_{i+1}.$$

(The indices j entering the union are finite in number due to Lemma 1.2.) Let λ_j be a sequence of positive numbers and let $\varphi = \sum_{j=1}^\infty \lambda_j \varphi_j$. Since $\{\text{supp } \varphi_j\}$ is locally finite, φ is finite as at any point only a finite number of terms give a non-vanishing contribution. Moreover, $\text{supp } \varphi = \bigcup_{j=1}^\infty \overline{U_j} \subset I^+(S_{-1})$ (it is useful to recall that the closure operator is additive if the union is over a locally finite family [9, Lemma 20.5], and that the closure of a locally finite family is locally finite).

We are going to find a sequence λ_j such that $\tau^- := \tau_\varphi^-$ is steep over $J^+(S_0)$. Observe that

$$\tau_\varphi^- = \sum_j \lambda_j \tau_{\varphi_j}^-, \quad \nabla \tau_\varphi^- = \sum_j \lambda_j \nabla \tau_{\varphi_j}^-.$$

By induction suppose that we can find a finite sequence $\lambda_1^i, \lambda_2^i, \dots, \lambda_{n_i}^i$, ($\lambda_j^i = 0$ for $j > n_i$) such that τ^- , defined as above, is steep over K_i , and satisfies the inequality $\tau^- - t > 0$ over K_i . Let Λ_i be the finite index set such that if $r \in \Lambda_i$ then $U_r \cap (K_{i+1} \setminus \text{Int } K_i) \neq \emptyset$. We have $U_r \cap K_{i-1} = \emptyset$, for otherwise $U_r \subset K_i \Rightarrow U_r \subset \text{Int } K_i$, a contradiction. Let $p \in K_{i+1} \setminus \text{Int } K_i$, then p belongs to some U_r , $r \in \Lambda_i$, and hence $E^-(p)$ intersects U_r .

We now replace the constants λ_r , $r \in \Lambda_i$, $\lambda_r \rightarrow \lambda'_r$, with larger constants chosen in

such a way that τ^- and $\nabla\tau^-$ get replaced by

$$\tau_\varphi^- \rightarrow \tau_\varphi^- + \sum_{r \in \Lambda_i} (\lambda'_r - \lambda_r) \tau_{\varphi_r}^-, \quad \nabla \tau_\varphi^- \rightarrow \nabla \tau_\varphi^- + \sum_{r \in \Lambda_i} (\lambda'_r - \lambda_r) \nabla \tau_{\varphi_r}^-.$$

and so that τ_φ^- becomes steep at p , and $\tau_\varphi^- - t$ becomes positive at p .

Since $K_{i+1} \setminus \text{Int} K_i$ is compact and $\tau^- - t$ and $\nabla\tau^-$ depend continuously on each λ_j , the redefinition of λ_r and consequently of τ^- can be done so as to obtain steepness and positivity of $\tau^- - t$ all over K_{i+1} . Observe that this redefinition can only increase the value of τ^- over K_i and that it certainly does not spoil steepness there. Furthermore, it does not change τ^- over K_{i-1} thus the built inductive process leads to the desired function τ^- . $\square$

Acknowledgments

I thank James Grant and Piotr Chruściel for useful and motivating discussions on this and related problems. This work has been partially supported by GNFM of INDAM.

- [1] A. N. Bernal and M. Sánchez. On smooth Cauchy hypersurfaces and Geroch's splitting theorem. *Commun. Math. Phys.*, 243:461–470, 2003.
- [2] P. T. Chruściel, J. D. E. Grant, and E. Minguzzi. On differentiability of volume time functions. *Ann. Henri Poincaré*, 2015. DOI:10.1007/s00023-015-0448-3, arXiv:1301.2909.
- [3] A. Fathi and A. Siconolfi. On smooth time functions. *Math. Proc. Camb. Phil. Soc.*, 152:303–339, 2012.
- [4] R. Geroch. Domain of dependence. *J. Math. Phys.*, 11:437–449, 1970.
- [5] S. W. Hawking and G. F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge University Press, Cambridge, 1973.
- [6] M. W. Hirsch. *Differential topology*. Springer-Verlag, New York, 1976.
- [7] O. Müller and M. Sánchez. Lorentzian manifolds isometrically embeddable in $\mathbb{L}^N$. *Trans. Am. Math. Soc.*, 363:5367–5379, 2011.
- [8] A. Rennie and B. E. Whale. Generalised time functions and finiteness of the Lorentzian distance. arXiv:1412.5652v1, 2014.
- [9] S. Willard. *General topology*. Addison-Wesley Publishing Company, Reading, 1970.